

AD-A091 293

GEORGIA INST OF TECH ATLANTA ENGINEERING EXPERIMENT --ETC F/G 20/5
A KINETIC STUDY OF THE REACTION OF CH₃ O₂ WITH NO₂. VOLUME I, (U)
1980 A R RAVISHANKARA, R L EISELE, P H WINE DOT-FA78WA-4259

UNCLASSIFIED

FAA/EE-80-38-VOL-1

NL

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

END
DATE
FILMED
12-80
DTIC

AD A091293

LEVEL II

A KINETIC STUDY OF THE REACTION OF CH_3O_2 WITH NO_2

by

A. R. Ravishankara

F. L. Eisele

P. H. Wine

12
A



DTIC
ELECTE
NOV 03 1980
S D E

DDC FILE COPY

Prepared for:

U.S. Department of Transportation
Contract No. DOT-FA-78WA-4259
Report No. FAA-EE-80-38
Volume I


GEORGIA INSTITUTE OF TECHNOLOGY

Engineering Experiment Station
Atlanta, Georgia 30332



DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited



80 10 30 004

18		19 80-38-VOL-1		Technical Report Documentation Page	
1. Report No. FAA/EE/80-35, vol. 1		2. Government Accession No. AD-A091293		3. Recipient's Catalog No. 11	
4. Title and Subtitle 6 A Kinetic Study of the Reaction of CH_3O_2 with NO_2 . Volume I,		5. Report Date 1980		6. Performing Organization Code	
7. Author(s) 10 A.R. Ravishankara, R.L. Eisele, P.H. Wine		8. Performing Organization Report No.		9. Performing Organization Name and Address Engineering Experiment Station Georgia Institute of Technology Atlanta, Georgia 30332	
10. Work Unit No. (TRAIS)		11. Contract or Grant No. 15 DOT-FA78WA-4259		12. Sponsoring Agency Name and Address U. S. Department of Transportation Federal Aviation Administration 800 Independence Ave., S. W. Washington, D. C. 20591	
13. Type of Report and Period Covered		14. Sponsoring Agency Code		15. Supplementary Notes 12/39	
16. Abstract					
17. Key Words			18. Distribution Statement This document is available to the public through the National Technical Information Service, Springfield, Va. 22161		
19. Security Classif. (of this report) Unclassified		20. Security Classif. (of this page) Unclassified		21. No. of Pages 38	
				22. Price 153850	

A KINETIC STUDY OF THE REACTION OF
 CH_3O_2 WITH NO_2

Contract #DOT-FA78WA-4259
Report # FAA-EE-80-38
VOLUME I

Submitted to:

U.S. Department of Transportation
Federal Aviation Administration
800 Independence Avenue, S.W.
Washington, D.C. 20591

by

Molecular Sciences Group
Electromagnetics Laboratory
Engineering Experiment Station
Georgia Institute of Technology
Atlanta, Georgia 30332

Accession For	
NTIS GAMA	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Mail and/or special
A	

Contracting through;

Georgia Tech Research Institute
Georgia Institute of Technology
Atlanta, Georgia 30332

Prepared by:

A.R. Ravishankara
F.L. Eisele
P.H. Wine

YIELD: AT K_{ant} 1

ABSTRACT


The technique of pulsed laser photolysis-long path laser absorption is employed to study the kinetics of the reaction $\text{CH}_3\text{O}_2 + \text{NO}_2 + \text{N}_2 \xrightarrow{k_1} \text{CH}_3\text{O}_2\text{NO}_2 + \text{N}_2$ over the temperature range 253-353 K. At all temperatures, k_1 is found to monotonically increase with increasing N_2 pressure over the range 76-722 torr, indicating that reaction occurs predominantly by addition. The dependence of k_1 on N_2 pressure shows that the reaction is in the fall-off region between second and third order kinetics. These results are compared with previous results and their atmospheric implications are discussed.

TABLE OF CONTENTS

	<u>Page</u>
INTRODUCTION	1
PULSED LASER PHOTOLYSIS-LONG PATH LASER ABSORPTION KINETICS STUDY OF THE REACTION OF METHYLPEROXY RADICALS WITH NO ₂	5
EXPERIMENTAL	6
RESULTS	11
DISCUSSION	22
REFERENCES	26
ACKNOWLEDGEMENTS	28
APPENDIX I	29
APPENDIX II	32

Introduction

The methylperoxy radical is an important intermediate in reaction sequences through which hydrocarbons are oxidized in both atmospheric and combustion processes. In combustion systems the fate of peroxy radicals is predominantly governed by their reactions with other radicals, while in the atmosphere, and particularly in the polluted troposphere, it is expected that reactions with stable molecules will be important. The major species which can react with CH_3O_2 in the atmosphere include HO_2 , NO , NO_2 , SO_2 , and O_3 . The rate constant for the reaction of CH_3O_2 with NO has been recently measured using direct kinetic techniques¹⁻³ and found to be nearly as fast as the reaction of HO_2 with NO .^{4,5} Therefore, in a polluted atmosphere where NO_x concentrations are high, methyl peroxy radicals are expected to react mostly with NO_x species. For the subject reaction of the present investigation,



to effectively compete with the reaction of CH_3O_2 with NO ,



the value of k_1 should be of similar magnitude as that of k_2 . If Reaction (1) were a significant path for CH_3O_2 removal, NO_2 production (and the subsequent O_3 generation) would be reduced.

Until recently most rate coefficients for CH_3O_2 radical reactions were obtained by steady state competitive experiments.⁶⁻⁸ In the past few years direct measurement techniques such as discharge flow-mass spectrometry, flash photolysis-ultraviolet absorption, and molecular

modulation-UV absorption spectroscopy have been adopted for CH_3O_2 radical studies.

To date three direct measurements of k_1 have been carried out. Cox and Tyndall⁹ utilized the molecular modulation-UV absorption technique and obtained a value of k_1 which was essentially independent of bath gas pressure. The flash photolysis-UV absorption method has been employed by Adachi and Basco,¹⁰ and Sander and Watson² to measure k_1 . Adachi and Basco's results are in agreement with those of Cox and Tyndall. However, unlike the previous two measurements, Sander and Watson found Reaction (1) to be pressure dependent. These investigators carried out their experiments under secondary reaction free conditions using a very sensitive probe for CH_3O_2 radicals. Hence, their results appear to be more reliable. There are no previous measurements of k_1 as a function of temperature.

In the present study, k_1 was measured as a function of N_2 diluent gas pressure and temperature using the recently developed technique of pulsed laser photolysis-long path laser absorption. Due to the monochromaticity of the photolysis laser, and the resultant photolysis specificity, all unwanted photofragments were essentially eliminated, while the high intensity of the CW 257 nm probing laser beam enabled measurement of very low concentrations of CH_3O_2 . Under these secondary reaction free conditions, our measured value of k_1 was found to be pressure dependent in excellent agreement with Sander and Watson. Also, as to be expected for an addition reaction, k_1 was found to increase with decreasing temperature.

The technique of pulsed laser photolysis-laser induced fluorescence was originally proposed to be employed in the study of Reaction (1). CH_3O_2 radicals were to be produced using either 355 nm photolysis of $\text{Cl}_2/\text{CH}_4/\text{O}_2$ mixtures or UV photolysis of azomethane. To follow the course of the reaction either NO_2 reactant in excess CH_3O_2 or CH_2O product in excess NO_2 were to be monitored using laser induced fluorescence.

There were two developments which excluded carrying out laser induced fluorescence detection of CH_2O to map out the course of Reaction (1): a) we could not obtain the expected 10 mW of UV laser output in the intracavity Ar^+ laser pumped tunable dye laser (in fact, we succeeded in obtaining only 20 μW !), and b) in some recent studies it was shown that Reaction (1) proceeds predominantly through an addition mechanism; therefore, the CH_2O product yield is very low, and an unrealistically high detection sensitivity for CH_2O would be needed. For these reasons, we abandoned the idea of monitoring CH_2O .

The second approach of following the NO_2 decay in excess CH_3O_2 was pursued when the first approach failed. The primary requirement for this experiment was the creation of a uniform known concentration of CH_3O_2 by laser photolysis. We have succeeded in obtaining a spatially uniform, 'top hat' profiled laser beam using a segmented aperture optical integrator. With this added capability, we can produce CH_3O_2 radicals in excess over NO_2 . Our capability to detect NO_2 via laser induced fluorescence (pumped by the 488 nm output of an argon ion laser) is well developed and has been used in the study of the reaction of CH_3O_2 with NO .³ However, when we tried to monitor NO_2 in excess CH_3O_2 , we found the decay of NO_2 to be non-exponential. The decay rates decreased with time due to the disproportionation reaction,



The rate coefficient for Reaction (a) is reasonably well known.² A simple calculation shows that unless Reaction (1) is very fast, i.e., is $\sim 1 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, the CH_3O_2 concentration would decrease by more than 10% by the time the NO_2 concentration dropped to $1/e$ times its original value. Therefore, it was evident that even though we could measure k_1 at low temperatures and high pressures the required range of pressures and temperatures could not be covered.

Upon realizing that our originally proposed methods would be either inapplicable or of limited use, we decided to develop the laser photolysis-long path laser absorption technique. This effort was very successful and we completed the measurement of k_1 as a function of temperature and pressure, as discussed below.

PULSED LASER PHOTOLYSIS-LONG PATH LASER ABSORPTION KINETICS
STUDY OF THE REACTION OF METHYLPEROXY RADICALS WITH NO₂

Experimental

A schematic diagram of the apparatus used to study Reaction (1) is shown in Figure 1. A jacketed reactor with an internal volume of $\sim 1\ell$ and a length of 107 cm was constructed out of pyrex. The cell was maintained at a constant temperature by circulating either methanol (253-298 K) or ethylene glycol (298-353 K) from a temperature controlled circulator through the outer jacket. It was found that the temperature was uniform to within 2°C throughout the length of the cell. On both ends of the reactor two $1/8$ " thick, 2" diameter quartz (Suprasil 1) plates were attached using O-ring seals and metal clamps. The reactants were pre-mixed in a mixing chamber and slowly flowed through the reactor. At the outlet of the reactor, the pressure in the system was measured using an absolute one-turn Bourdon gauge (Wallace and Tiernan Model FA160-NN) before the gas mixture was pumped out.

Azomethane/ N_2 and NO_2/N_2 mixtures were prepared in 12 ℓ bulbs on a gas handling system. The bulbs were connected to the mixing chamber using teflon tubing and stainless steel ultra-torr fittings. The rates of flow of the azomethane/ N_2 mixture, NO_2/N_2 mixture, N_2 , and O_2 into the mixing chamber were controlled by stainless steel needle valves and measured by calibrated mass flow transducers. The tubes leading into the mixing chamber had radial outlets to turbulently mix the gases. The concentration of each component in the reaction mixture was determined from measurements of appropriate mass flow rates and the total pressure. The concentration of NO_2 in the NO_2 /diluent mixture was checked (before and after each set of experiments) by simultaneous measurements of the total pressure (of the mixture) and NO_2 absorption




Figure 1. A Schematic Diagram of Pulsed Laser Photolysis-Long Path Laser Absorption Apparatus.

at 366 nm. The measurements were carried out using an Hg pen-ray lamp as the light source, a 70 cm absorption cell, and a bandpass filter-photo-multiplier detector combination. The absorption cross section used to determine $[\text{NO}_2]$, $5.75 \times 10^{-10} \text{ cm}^2$,¹¹ has been previously measured in our laboratory and agrees well with the literature value.¹² The azomethane concentration was occasionally checked by a method identical to that for NO_2 ; the absorption cross section of azomethane was measured to be $1.13 \times 10^{-20} \text{ cm}^2$ at 366 nm, which is slightly higher than the published value,¹³ but in agreement with other recent measurements.¹⁴

Azomethane was prepared as described in Appendix I. NO_2 , obtained from Matheson Gas Products, had a stated purity of 99.5%, and was purified by mixing it with UHP O_2 to convert all NO to NO_2 , and subsequently subjecting it to freeze-pump-thaw cycles to remove O_2 . NO_2 and O_2 were Matheson grade and had stated purity levels of 99.9995% and 99.99%, respectively; these gases were used as supplied.

An ArF excimer laser, the photolysis light source, was housed in an adjoining room, approximately 2 meters from the front window of the reactor. We found this spatial separation to be essential if a uniform beam was to be obtained. (Operation of the laser under slightly fluorine-rich conditions also helped to make the beam more diffuse.) The beam was approximately 2 cm x 7 cm by the time it reached the reactor. The laser beam was transmitted through a custom made dielectric mirror, DM and traveled through the reactor impinging on the MgF_2 coated aluminum mirror M_4 where a fraction of the beam was reflected back into the reactor. Both mirrors DM and M_4 were mounted on precision mirror mounts with vernier control to enable very fine adjustments. (DM was specially coated for us by Acton Research Corp. to have $\sim 85\%$ transmission at 193 nm and $\geq 95\%$ reflectivity at 257 nm.)

The doubled CW 257 nm UV beam was generated as described in Appendix II. This beam was diverted into the reactor using mirror M_1 so as to traverse the volume photolyzed by the ArF laser beam. By adjusting Mirrors M_1 and M_4 , the CW UV beam was multipassed 8 times through the reactor to obtain a path length of ~ 860 cm. The 8th pass was diverted away from the reactor by a small aluminum mirror M_2 , passed through a methanol cell and a 257 nm bandpass filter, and detected by a quartz envelope 1P28 (PQA) photomultiplier tube. (It was necessary to use a methanol filter to prevent the 193 nm laser pulse from being detected by the PM tube.) The output of the PM tube was amplified and stored in a signal averager (Tracor-Northern 1500) operating in the analog mode and triggered 150 μ sec after the photolysis pulse. (We could not obtain a usable signal in the first 150 μ sec due to interference from fluorescence induced by the 193 nm beam.) The PM tube was operated at ~ 600 V to have a resistor bridge current of ~ 1 mA. The anode current was always kept below 1 μ A by using suitable neutral density filters to reduce the intensity of the 257 nm beam. To increase the signal-to-noise ratio of the decay curves, 16 to 64 flashes were averaged. The pulse rate of the photolysis laser (0.03 to 0.01 Hz) was such that the contents of the reactor was replaced between consecutive photolysis pulses. This was necessary in order to sweep out the products (which also absorb the probing beam) and use identical gas mixtures for each flash. In preliminary experiments it was established that as long as the cell was swept out between flashes either the laser repetition rate or the flow rate of the mixture through the reactor did not affect the measured value of k_1 . The intensity of the transmitted 257 nm beam before photolysis was higher than that obtained when all CH_3O_2 had reacted. This residual absorption has been observed by both Cox and

Tyndall,⁹ and Sander and Watson,² and is attributed to the absorption by $\text{CH}_3\text{O}_2\text{NO}_2$, the product of Reaction (1). Sander and Watson² have shown that the absorption of the analyzing beam by a reaction product which is stoichiometrically coupled to the monitored reactant species by a single reaction (as it is in the present case) does not affect the kinetic data that is obtained. In addition, they point out that the proper I_0 value that should be used in the data analysis is the intensity of the 257 nm beam when all CH_3O_2 has reacted, and not the value obtained when $\text{CH}_3\text{O}_2\text{NO}_2$ is absent. It is worth noting that since the intensity of the probing beam was high, the signal-to-noise was excellent and we could easily work with less than 3% initial absorption and still follow the decay of CH_3O_2 down to ~ 4 $1/e$ times!

Results

Reaction (1) was studied as a function of the pressure of the bath gas N_2 and temperature (at 298K, 76-722 torr; at 353K, 330-696 torr; at 253K, 109-510 torr). As pointed out earlier, all experiments were carried under pseudo-first order conditions with $[NO_2] > [CH_3O_2]$. Figure 2 shows typical decays of $[CH_3O_2]$ as a function of time; the decays are exponential and thus confirm the presence of pseudo-first order conditions. The decay of $[CH_3O_2]$ was usually followed, for 3 $1/e$ times. The pseudo-first order rate constant, k_1' (\equiv slope of $\ln[CH_3O_2]$ vs. t plots) was measured for various values of $[NO_2]$ at each temperature and N_2 pressure. k_1' was then plotted against $[NO_2]$; the slope of this line gave k_1 . Figure 3 shows one such plot. It should be pointed out that the fitting procedure we employed (a linear least squares analysis) did not force the k_1' vs. $[NO_2]$ line to a zero intercept; in our system, it is possible to have a non-zero intercept since the entire volume of the reactor was not subjected to the photolysis. The intercepts in all our experiments were small relative to the measured k_1' values.

Table I lists the individual values of k_1' along with all the pertinent experimental conditions. The quoted errors for k_1 , calculated from the slope of k_1' vs. $[NO_2]$ plots, are 2σ and reflect the precision of the measurement. The absolute accuracy of the measured values of k_1 , is expected to be $\sim 20\%$ which includes the error in the $[NO_2]$ measurement.

Table II lists the values of k_1 obtained at various N_2 pressures and temperatures. Figure 4 shows a plot of k_1 as a function N_2 number density at three different temperatures. The solid lines drawn through these points are visual best fits.




Figure 2. $[\text{CH}_3\text{O}_2]$ Temporal Profiles in the Presence of Varying Amounts of NO_2 . The concentration of NO_2 for each decay curve is shown next to the least squares lines drawn through the experimental points.




Figure 3. Plot of Pseudo-First Order Rate Constants vs. NO_2 Concentration at 298K and 519 torr of N_2 .

Table I. Rate Constant Data for the Reaction of CH_3O_2 with NO_2 .

Temperature K	N_2 Pressure torr	[Azomethane] 10^{15} cm^{-3}	$[\text{O}_2]$ 10^{17} cm^{-3}	$[\text{CH}_3\text{O}_2]_0^a$ 10^{12} cm^{-3}	$[\text{NO}_2]$ 10^{14} cm^{-3}	$k_1, 10^3 \text{ s}^{-1}$	$k_1 \times 10^{12b}$ $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$
298	76±3	0.80	2.70	2.29	6.95	1.15	
		0.78	2.69	1.96	11.73	1.75	
		0.74	2.60	1.31	19.10	2.56	
		0.70	2.63	0.90	24.81	3.64	(1.36±0.23)
	157±2	0.52	4.50	1.13	7.94	1.50	
		1.52	4.34	1.22	10.08	1.70	
		0.53	4.34	1.00	14.96	2.82	
		0.81	4.40	2.33	17.54	3.12	
		0.53	4.47	0.99	21.47	4.05	(1.90±0.32)
	258±3	0.82	3.46	5.60	5.65	1.39	
		0.85	3.50	7.02	12.26	2.90	
		0.79	3.42	6.91	15.60	4.04	
		0.79	3.54	7.00	19.8	5.60	(2.60±0.56)
	352±3	0.77	3.53	2.56	6.87	1.88	
		0.71	3.52	3.39	9.42	2.70	
		0.79	3.52	2.46	11.88	3.70	
		0.67	3.50	2.32	13.63	3.77	
		0.73	3.51	2.66	16.05	4.50	(2.82±0.28)
	519±2	1.14	3.82	2.17	6.21	1.84	
		1.15	3.82	2.85	11.70	3.63	
		1.16	3.78	3.28	13.59	4.57	
		1.23	3.79	3.46	15.95	4.93	
		1.08	3.80	4.80	21.60	7.06	(3.36±0.32)
	722±1	0.821	5.68	3.75	5.38	1.91	
		1.76	5.66	4.00	7.47	2.60	
		0.843	5.68	4.70	9.86	3.42	
		0.88	5.66	5.90	12.79	4.95	
		0.91	5.66	3.20	15.92	6.15	(4.12±0.38)

Temperature K	N ₂ Pressure torr	[Azomethane] 10 ¹⁵ cm ⁻³	[O ₂] 10 ¹⁷ cm ⁻³	[CH ₃ O ₂] 10 ¹³ cm ⁻³	[NO ₂] 10 ¹⁴ cm ⁻³	k _f , 10 ³ s ⁻¹	k ₁ × 10 ^{12b} cm ³ molecule ⁻¹ s ⁻¹
353	330±3	0.92	3.02	2.98	4.60	0.70	
		0.91	3.50	3.30	9.50	1.38	
		0.93	2.50	1.39	11.81	1.61	
		0.87	2.50	1.70	15.83	2.09	(1.19±0.15)
	354±2	0.96	4.57	2.22	8.64	1.42	
		0.86	4.56	1.97	15.32	2.27	
		1.03	4.56	1.35	24.73	3.81	
		0.79	4.53	1.03	30.21	4.22	(1.35±0.21)
	511±2	1.13	4.18	2.46	7.81	1.78	
		1.36	4.17	3.89	17.44	3.34	
		1.21	4.19	2.18	23.60	4.37	
		1.22	4.16	2.65	28.89	5.41	(1.71±0.10)
253	696±2	1.11	4.76	2.23	9.94	2.28	
		2.08	4.69	3.57	20.70	4.17	
		1.74	5.00	3.12	24.12	5.27	
		1.46	4.73	1.71	31.71	6.81	
	109±1	1.62	4.73	2.41	4.50	8.84	(1.93±0.20)
		1.30	3.60	1.53	2.77	1.05	
		1.39	4.56	1.55	4.02	1.44	
		1.32	3.60	1.62	5.75	2.03	
	250±2	1.31	3.70	1.82	7.56	2.36	
		1.31	3.80	0.89	9.16	2.86	
		1.31	3.70	1.31	11.10	3.10	(2.50±0.34)
		0.57	2.23	1.40	2.61	1.17	
		0.59	2.22	1.62	3.42	1.38	
		0.61	2.22	1.32	5.66	2.29	
		0.61	2.22	1.54	7.89	3.29	
		0.64	2.21	1.60	9.96	3.89	(3.85±0.30)

Temperature K	N ₂ Pressure torr	[Azomethane] 10 ¹⁵ cm ⁻³	[O ₂] 10 ¹⁷ cm ⁻³	[CH ₃ O ₂] ₀ ^a 10 ¹³ cm ⁻³	[NO ₂] 10 ¹⁴ cm ⁻³	k ₁ , 10 ³ s ⁻¹	k ₁ × 10 ¹² ^b cm ³ molecule ⁻¹ s ⁻¹
253	503±3	0.60	4.59	1.80	3.00	1.66	
		0.78	5.36	1.30	4.34	2.16	
		0.63	4.56	2.55	6.55	3.01	
		0.62	4.59	2.59	8.52	4.05	(5.10±0.50)
519	519	1.11	5.22	2.2	5.85	2.60	
		0.99	5.21	2.6	11.20	5.72	
		0.88	5.21	2.0	18.15	9.72	
		1.10	5.22	2.1	23.8	13.20	(5.8±1.0)

a) The initial concentration of CH₃O₂, [CH₃O₂]₀, was calculated by back-extrapolating the decay curve for 150 μsec, (the delay between photolysis and initiation of signal acquisition.)

b) The quoted errors are 2σ and represent the precision of the measurements.

Table II. Rate Constants for the Reaction of CH_3O_2 with NO_2 as a Function of Temperature and N_2 Pressure.

Temperature K	Pressure torr	Number Density $10^{19}\text{molecules cm}^{-3}$	$k_1 \times 10^{12}$ $\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$
298	76	0.246	1.36 ± 0.23
	157	0.509	1.90 ± 0.32
	258	0.836	2.60 ± 0.56
	352	1.141	2.82 ± 0.28
	500	1.620	3.25 ± 0.50
	519	1.682	3.36 ± 0.32
	722	2.339	4.12 ± 0.38
353	330	0.903	1.19 ± 0.15
	354	0.966	1.35 ± 0.21
	511	1.398	1.71 ± 0.10
	696	1.904	1.93 ± 0.20
253	109	0.416	2.50 ± 0.34
	250	0.954	3.85 ± 0.30
	503	1.920	5.10 ± 0.50
	519	1.981	5.8 ± 1.0




Figure 4. Plots of k_1 vs. N_2 Number Density. The lines drawn through the points are visual best fits.

There are two possible chemical complications that can arise in our experiments: (a) production of O_3 due to reaction of $O(^3P)$ (formed by either O_2 or NO_2 photolysis) with O_2 and (b) production of NO and its subsequent reaction with CH_3O_2 . Photolysis of O_2 under the most unfavorable conditions, i.e., high $[O_2](\sim 5 \times 10^{17} \text{ cm}^{-3})$ and high photolysis laser energy ($\sim 10 \text{ mJ/cm}^2$), would produce $\sim 1 \times 10^{13} O(^3P) \text{ cm}^{-3}$. (The absorption cross section for O_2 at 193 nm is $\sim 1 \times 10^{-21} \text{ cm}^2$.¹³) Similarly, photolysis of NO_2 would lead to $\sim 1 \times 10^{13} O(^3P) \text{ cm}^{-3}$. (The absorption cross section for NO_2 at 193 nm is $\sim 6 \times 10^{-19} \text{ cm}^2$.¹⁵) Any $O(^1D)$ that is formed via photolysis would be quenched to $O(^3P)$ in less than 0.1 μsec by N_2 .¹⁶ The photolytically produced $O(^3P)$ would then react with either O_2 or NO_2 ,



The fraction of $O(^3P)$ that would lead to O_3 formation would, of course, depend on the ratio of $k_4[O_2][M]$ to $k_5[NO_2]$. Under the extreme conditions of low temperature (where k_4 is large) and low NO_2 , we would produce $\sim 4 \times 10^{12} O_3 \text{ cm}^{-3}$. In our experiment, then, we would obtain $\sim 3\%$ absorption of 257 nm radiation due to O_3 . However, the O_3 formation does not affect measurement of the CH_3O_2 decay since all O_3 is formed within 100 μsec (99% completion) after the laser flash, and, therefore simply changes the value of I_0 . The concentration of O_3 formed is too low to have any effect on the kinetics of the measured rate of CH_3O_2 decay. This conclusion was checked by increasing the photolysis energy by a factor of 4 while keeping $[CH_3O_2]_0$ constant (by decreasing [azomethane] by a factor

of 4); the measured decay rates were, within experimental error, unaffected.

The amount of NO that is produced via NO₂ photolysis would be at most 1.2% when 10 mJ/cm² is used for photolysis and all O(³P) that is produced reacts with NO₂ to produce NO. NO would of course, react with CH₃O₂ to produce CH₃O and NO₂,



The rate constant for Reaction (2) is known to be $\sim 8 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.¹⁻³ Since the measured values of k_1 vary from 1.19 to $5.8 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, the contribution of Reaction (2) could never be greater than 8% (i.e., at high temperature and low pressures). Again, this possibility was discarded as inconsequential by the observation that the measured rate constant was independent of photolysis energy. CH₃O formed in Reaction (2) would react with NO₂ to give CH₃ONO₂.



Formation of CH₃ONO₂ would not affect the measured rate constant since CH₃ONO₂ does not regenerate CH₃O₂, does not strongly absorb 257 nm radiation, and its concentration is too low to effectively remove CH₃O₂ via reaction. (The absorption cross section for CH₃ONO₂ is $\sim 4 \times 10^{-20} \text{ cm}^2$ at 257 nm.¹⁷)

In addition to the two photolysis related problems that were discussed above, it is possible that our measured rate constants are incorrect if the thermal decomposition of the CH₃O₂NO₂ product leads to CH₃O₂ production on the same time scale as the measured CH₃O₂ decay,



Reaction (7) would, of course, be most severe at higher temperatures. The rate constant for Reaction (7) is not known. However, it cannot be very different from that of HO_2NO_2 decomposition leading to HO_2 and NO_2 production, which has been measured.¹⁸ Using this rate constant, the fastest rate for Reaction (7) at 353K, would be $\sim 20\text{s}^{-1}$. This rate is negligible when compared to the $[\text{CH}_3\text{O}_2]$ decay rates of 699-8837 s^{-1} that were measured at 353K. Experimentally, if Reaction (7) was important, we would have observed non-exponential $[\text{CH}_3\text{O}_2]$ decays and more importantly, a non-linear dependence of k_1' on $[\text{NO}_2]$. We found the $[\text{CH}_3\text{O}_2]$ to decay exponentially for at least 3 $1/e$ times and k_1' vs $[\text{NO}_2]$ plots were linear. Therefore, Reaction (7) could not have been significant under our experimental conditions.

Discussion

There are three separate direct measurements of k_1 at 298K. Cox and Tyndall⁹ measured k_1 to be $(1.2 \pm 0.3) \times 10^{-12} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ at 500 torr of Ar + CH₄, and $k_1 = (1.6 \pm 0.3) \times 10^{-12} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ at 540 torr N₂, using the molecular modulation technique. This pressure independent value of k_1 is in good agreement with those obtained by Adachi and Basco¹⁰ who measured k_1 to be $(1.53 \pm 0.07) \times 10^{-12} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ independent of pressure (between 53 and 580 torr Ar) using the method of flash photolysis-kinetic spectroscopy. This pressure independent behavior has been interpreted to mean that Reaction (1) is at its high pressure limit above ~ 50 torr of Ar. Sander and Watson,² however, found k_1 to vary from $(1.5 \pm 0.10) \times 10^{-12} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ at 50 torr N₂ to $(3.94 \pm 0.17) \times 10^{-12} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ at 700 torr of N₂ at 298K. Moreover, the value of k_1 was observed to be dependent on the nature of the bath gas, i.e., He, N₂, or SF₆. This type of pressure dependent behavior is typical of an addition reaction. Sander and Watson's data also shows that Reaction (1) is in the "fall-off" regime, i.e., between second and third order, in pressure range of 50 to 700 torr of He, N₂, and SF₆. It should also be pointed out that the lower pressure values of k_1 obtained by Sander and Watson agree reasonably well with k_1 values of both Cox and Tyndall, and Adachi and Basco.

As seen from Table III, our measured values of k_1 are in excellent agreement with those of Sander and Watson; this argues very heavily for k_1 being pressure dependent. Sander and Watson have pointed out that the possible reason for the discrepancy between their values at higher pressures and those of Cox and Tyndall, is the long residence time of the gas mixtures and depletion of NO₂ along the length of the cell in

Table III. Comparison of Our Results With Previous Data at 298K.

Pressure torr	Diluent	This Work	$k_1, 10^{12} \times \text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$		
			Sander and Watson Ref (2)	Cox and Tyndall Ref (9)	Adachi and Basco Ref (10)
50	N ₂		1.15±0.10		
76	N ₂	1.36±0.23			
100	N ₂		1.58±0.15		
157	N ₂	1.90±0.32			
225	N ₂		2.22±0.31		
258	N ₂	2.60±0.56			
350	N ₂		2.98±0.23		
352	N ₂	2.82±0.28			
500	N ₂	3.25±0.50	3.67±0.21		
519	N ₂	3.36±0.32			
540	N ₂			1.6±0.3	
700	N ₂		3.94 ± 0.17		
722	N ₂	4.12±0.38			
50	Ar+CH ₄			1.2±0.3	
53-580	Ar				1.53±0.07

* The preferred value reported by Watson and Sander is the average value $\equiv k_1' / [\text{NO}_2]$.

the molecular modulation apparatus. In Adachi and Basco's experiments, the concentrations of reactants used were extremely large. This should have led to many secondary reaction complications. Also, their CH_3O_2 radical detection scheme was quite insensitive. Furthermore, as the authors pointed out, they would have to make a large correction to their high flash energy results due to the participation of Reaction (6), if $k_6 \approx 8 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. To date, three separate direct measurements of k_6 have been carried out which yield a value of $k_6 = (8 \pm 2) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.¹⁻³ Even though Adachi and Basco carried out a separate set of experiments using low flash energies and obtained values of k_1 (supposedly in the absence of secondary reaction complications) in agreement with their previous measurements, their results have to be considered suspect. They had very large concentrations of free radicals even in their low flash energy experiments which could lead to erroneous results. We believe that at 298K, the data obtained by Sander and Watson and in the present investigation are devoid of secondary reaction complications and hence should be used for atmospheric modeling studies.

There are no previous determinations of the temperature dependence of k_1 . Our results are consistent with what is to be expected for an addition reaction which goes through an energy-rich complex. The strong temperature dependence of the measured values of k_1 suggests that $k_{1\infty}$ is likely to be $> 1 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 253K. At 600 Torr, for example, the apparent "activation energy" for Reaction (6) is $\sim -1.8 \text{ Kcal/mole}$.

For atmospheric modelling purposes it is convenient to obtain an analytical expression which best describes the temperature and pressure dependence of k_1 . Sander and Watson have fit their data for k_1 at 298K

to the following semi-empirical equation which is based on Troe's formulation of RRKM theory:¹⁹

$$k([M], T) = \frac{k_0(T)[M]}{1 + \frac{k_0(T)[M]}{k_\infty(T)}} F_c \{1 + [\log_{10}(k_0(T)[M]/k_\infty(T))]^2\}^{-1} \quad (I)$$

$$k_0(T) = k_0(298) \left(\frac{T}{298}\right)^{-n}$$

$$k_\infty(T) = k_\infty(298) \left(\frac{T}{298}\right)^{-m}$$

In Equation (I), k_0 (k_∞) are the rate constants in the low (high) pressure limits and F_c is a parameter which relates the energy dependence of the rate constant for activated complex decomposition to the shape of the k vs $[M]$ curve. They obtain $k_0(298) = (2.33 \pm 0.8) \times 10^{-30} \text{ cm}^6 \text{ molecule}^{-1} \text{ sec}^{-1}$, $k_\infty = (7.0 \pm 1.0) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ sec}^{-1}$ and $F_c = 0.4$. Because Sander and Watson's data base at 298K is much larger than ours and because our results are in excellent agreement with theirs, we have fit our temperature dependent data by trial and error with the constraint that $k_0(298)$, $k_\infty(298)$, and F_c must fall within the error limits given by Sander and Watson. The best fit to the data is obtained for (Figure 4)

$$k_0(T) = 2.2 \times 10^{-30} \left(\frac{T}{298}\right)^{-2.5} \text{ cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$$

$$k_\infty(T) = 7 \times 10^{-12} \left(\frac{T}{298}\right)^{-3.5} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$$

$$F_c = 0.4$$

Our results suggest that Reaction (1) could contribute significantly towards removal of CH_3O_2 and NO_2 in the presence of high concentrations of NO_x (such as those encountered in urban smog) if $\text{CH}_3\text{O}_2\text{NO}_2$ is thermally stable.

References

1. I.C. Plumb, K.R. Ryan, J.R. Steven, and M.F.R. Mulcahy, Chem. Phys. Letters 63, 255 (1979).
2. S.P. Sander and R.T. Watson, J. Phys. Chem. (in press).
3. A.R. Ravishankara, F.L. Eisele, and P.H. Wine, presented at the Gordon Conference on Environmental Chemistry—Air, 1979.
- 4.(a) C.J. Howard and K.M. Evenson, Geophys. Res. Lett. 4, 437 (1977).
(b) B.Reimann and F. Kaufman, paper presented at 13th Informal Conference on Photochem., Clearwater Beach, FL (1978).
(c) J.P. Burrows, D.I. Clift, G.W. Harris, B.A. Thrush, and J.P. Wilkinson, W.M.O. Sym. Ozone, Toronto, Canada (1978).
(d) J.J. Margitan and J.G. Anderson, paper presented at 13th Informal Conference on Photochemistry, Clearwater Beach, FL (1978).
(e) M.T. Leu, J. Chem. Phys. 70, 1662 (1979).
5. C.J. Howard, J. Chem. Phys. 71, 2352 (1979).
- 6.(a) C.W. Spicer, A. Villa, H.A. Wiebe, and J. Heicklen, J. Amer. Chem. Soc. 95, 13 (1973).
(b) R. Simonaitis and J. Heicklen, J. Phys. Chem. 78, 2417 (1974).
(c) R. Simonaitis and J. Heicklen, J. Phys. Chem. 79, 298 (1975).
7. C.T. Pate, B.J. Finlyason, and J.N. Pitts, Jr., J. Amer. Chem. Soc. 96, 6554 (1974).
8. R.A. Cox, R.G. Derwent, P.M. Holt, and J.A. Kerr, J.C.S. Faraday I, 72, 2444 (1976).
9. R.A. Cox and G.S. Tyndall, Chem. Phys. Lett. 65, 357 (1979).
10. H. Adachi and N. Basco. Int. J. Chem. Kinet. XII, 1 (1980).
11. P.H. Wine, N.M. Kreutter, and A.R. Ravishankara, J. Phys. Chem. 83, 3191 (1979).
12. A.M. Bass, A.E. Ledford, Jr., and A.H. Laufer, J. Res. Natl. Bur. Stand. Section A 80, 143 (1976).
13. J.G. Calvert and J.N. Pitts, Jr., "Photochemistry," John Wiley and Son, Inc. 1966.
14. S.P. Sander and R.T. Watson, Private communications, 1980.
15. H.S. Johnston, S.G. Chang, and G. Whitten 78, 1 (1974).

16. NASA Reference Publication 1049, "The Stratosphere: Present and Future," Editors, R.D. Hudson and E.I. Reed, (1979).
17. W.D. Taulor, T.D. Allston, M.J. Moscato, G.B. Fazekas, R. Kozlosi, and G.A. Takacs, Int. J. Chemical Kinetics, in press, 1980, and references therein.
18. R.A. Graham, A.M. Winer, and J.N. Potts, Jr., J. Chem. Phys. 68, 4505 (1978).
19. J. Troe, J. Phys. Chem. 83, 114 (1979).
J. Troe, J. Chem. Phys. 66, 4758 (1977).

Acknowledgements

We would like to thank Drs. S.P. Sander and R.T. Watson for very helpful discussions throughout this work. We would also like to thank Mr. A.O. Langford for discussions which steered us towards utilization of colinear photolysis and probing laser beams.

This work was supported by the U.S. Department of Transportation/ Federal Aviation Administration through Contract #DOT-FA78WA-4259.

APPENDIX I

Preparation of Azomethane

The apparatus used for preparation of azomethane is shown in Figure 1a. The method we used is a slightly modified version of the Renaud and Leitch preparation procedure.

Ten grams of Sym-1,2-dimethylhydrazinedihydrochloride (obtained from Aldrich Chemical Co.) was slowly added to 25 ml of 6N NaOH contained in a 250 ml Erlenmeyer flask. The flask was cooled in an ice bath and the contents vigorously stirred using a magnetic stirrer. When 10 gm of Sym 1,2-dimethylhydrazinedihydrochloride was added, ~ 3 ml of 6N NaOH was mixed with this solution to dissolve any remaining solid. This solution was transferred to the dropping funnel.

35 grams of mercury (II) oxide was mixed with 50 ml of distilled water in the three-necked 500 ml flat-bottom flask. The three-necked flask was connected on one side to a helium tank and another side to a cold finger. The cold finger, maintained at 0°C with crushed ice, was connected to two glass traps in series. Both traps were cleaned by flaming them under vacuum.

Before starting the addition of the solution present in the dropping funnel to the three-necked flask, the assembled system was flushed with He for at least one hour to remove all traces of air present in the assembly. The first trap was maintained at 193K (dry ice—trichloroethylene slush) and the second at ~ 163K (ethanol cooled with liquid N₂). The solution in the dropping funnel was added in drops to the slurry of mercuric oxide which was slowly stirred with a magnetic stirrer. The addition was completed in ~ 45 minutes and He allowed to flow for another




Figure 1a. Apparatus for the Preparation of Azomethane.

20 minutes to sweep out any remaining azomethane in to the traps. (The flow rate of helium was ~ 20 ml/min.)

The traps were disconnected from the assembly and cooled to 77K with liquid N_2 to pump out He before transferring the contents to a bulb. The cold finger of the bulb was then maintained at 197K and azomethane distilled into a storage vessel maintained at 77K. The first and the last fractions were discarded. The vapor pressure of azomethane at 197K is ~ 7 torr while that of H_2O is less than 1×10^{-3} torr which enabled us to separate the two.

All preparation and handling of azomethane was carried out in the dark or under red lights to avoid photolysis. The azomethane was stored at 197K.

APPENDIX II

Generation of the 257 nm CW UV Laser Beam

A schematic diagram of the setup is shown in Figure 2a. A Lexel model 95 Argon ion laser was operated on the 514 nm line. The output of this laser was checked to ensure that the laser was operating in TEM₀₀ mode. The Gaussian beam was focused using a 10 cm focal length quartz plano convex lens to obtain a tight beam waist with a confocal parameter of 10 cm. (The focal length at 514 nm of the lens was accurately measured using the cat's eye technique) The position of the lens was adjusted to locate the beam waist in the middle of a 50 mm long (10 mm x 10 mm on the side) ADP crystal (Inrad Corp.) which was housed in a gold plated copper block. This copper block was in snug contact with a gold plated pedestal which was backed by a Peltier thermoelectric cooler. The Peltier cooler itself was backed by a heater block which, in turn, was cooled by circulating water at a constant temperature ($\pm 2^\circ\text{C}$) from a temperature controlled five gallon water bath heated by a fish tank heater. By adjusting the rate of water flow and the temperature of the heater block (which was controlled by a proportional heater) the temperature of the ADP crystal could be maintained at -10°C with a precision of $\pm 0.1^\circ\text{C}$. We found that the doubled 257 nm output was a maximum at this temperature.

The 257 nm beam was diverging as it exited the crystal; this beam was rendered parallel by a second lens (~ 15 cm focal length at 257 nm). The collinear 257 nm and 514 nm beams were dispersed by a prism and the 514 nm beam stopped by a block. The entire assembly was housed inside a box to prevent dust from getting in and 514 nm light from scattering out.




Figure 2a. Details of the Second Harmonic Generation Set Up.

With an input energy of 1 W at 514 nm, we could routinely obtain
~ 50 μ W of 257 nm, which was more than adequate to carry out our experiments.

2-8